

Contribution of groundwater carbon pools to atmospheric carbon sinks: A case study of the Yinchuan Basin, Northwest China

Hong-yun Ma, Jiang Liu, Hong-na Ma, Hua-qi Wang, Cheng-zhu Li, Yan-e Li

Citation: Hong-yun Ma, Jiang Liu, Hong-na Ma, Hua-qi Wang, Cheng-zhu Li, Yan-e Li, 2025. Contribution of groundwater carbon pools to atmospheric carbon sinks: A case study of the Yinchuan Basin, Northwest China, China Geology, 8, 295-302. doi: 10.31035/cg20230102.

View online: https://doi.org/10.31035/cg20230102

Related articles that may interest you

Global significance of the carbon cycle in the karst dynamic system: evidence from geological and ecological processes

China Geology. 2018, 1(1), 17 https://doi.org/10.31035/cg2018004

Carbon peak and carbon neutrality in China: Goals, implementation path and prospects

China Geology. 2021, 4(4), 720 https://doi.org/10.31035/cg2021083

Sedimentology and sequence stratigraphy of the mixed clastic-carbonate deposits in the Late Paleozoic icehouse period: A case study from the northern Qaidam Basin

China Geology. 2021, 4(4), 673 https://doi.org/10.31035/cg2021068

Major contribution to carbon neutrality by China's geosciences and geological technologies

China Geology. 2021, 4(2), 329 https://doi.org/10.31035/cg2021037

Origin and depositional environments of source rocks and crude oils from Niger Delta Basin: Carbon isotopic evidence

China Geology. 2020, 3(4), 602 https://doi.org/10.31035/cg2020057

A fast identification method based on the typical geophysical differences between submarine shallow carbonates and hydrate bearing sediments in the northern South China Sea

China Geology. 2020, 3(1), 16 https://doi.org/10.31035/cg2020021

China Geology 8 (2025) 295−302

China Geology

Journal homepage: http://chinageology.cgs.cn
https://www.sciencedirect.com/journal/china-geology

Contribution of groundwater carbon pools to atmospheric carbon sinks: A case study of the Yinchuan Basin, Northwest China

Hong-yun Ma[a, b, c], Jiang Liu[a, b,*], Hong-na Ma[a, b], Hua-qi Wang[a, b, c], Cheng-zhu Li[a, b], Yan-e Li[a, b]

[a] *Xi'an Center, China Geological Survey, Ministry of Natural Resources, Xi'an* 710054, *China*
[b] *Northwest China Center for Geoscience Innovation, China Geological Survey, Ministry of Natural Resources, Xi'an* 710054, *China*
[c] *Key Laboratory for Groundwater and Ecology in Arid and Semi-arid Regions, China Geological Survey, Ministry of Natural Resources, Xi'an* 710054, *China*

ARTICLE INFO

Article history:
Received 6 September 2023
Received in revised form 31 January 2024
Accepted 27 February 2024
Available online 27 March 2024

Keywords:
Carbon sink
Groundwater carbon pool
Surface soil carbon pool
Hydrogeochemistry
Carbon migration
Irrigation basin
Carbon burial engineering
Yinchuan Basin
Northwest China

ABSTRACT

Addressing global warming, a common change today, requires achieving peak carbon dioxide emissions and carbon neutrality (also referred to as the dual carbon goals). Enhancing research on the carbon cycle is urgently needed as the foundation. Water, a key carrier in the carbon cycle, necessitates investigation into groundwater carbon pools' contribution to atmospheric carbon sinks. This study assessed carbon stocks in the Yinchuan Basin's soil and groundwater carbon pools. Findings indicate the basin's surface soils contain approximately 24.16 Tg of organic carbon and a total of 60.01 Tg of carbon. In contrast, the basin's groundwater holds around 4.90 Tg of carbon, roughly one-fifth of the organic carbon in surface soils. Thus, groundwater and soil carbon pools possess comparable carbon stocks, underscoring the importance of the groundwater carbon pool. Studies on terrestrial carbon balance should incorporate groundwater carbon pools, which deserve increased focus. Evaluating groundwater carbon pools' contributions is vital for achieving the dual carbon goals.

©2025 China Geology Editorial Office.

1. Introduction

To proactively address global climate change and mitigate natural disasters like extreme weather, China has initiated the due carbon goals and has been actively engaged in research on the carbon cycle. Such research highlights the investigation of the dynamics of the carbon cycle and carbon budget of terrestrial ecosystems, as minor alterations in these ecosystems' carbon pools can cause considerable shifts in atmospheric CO_2 concentration (Piao SL et al., 2019; Wang K et al., 2023).

Terrestrial ecosystem carbon primarily resides in two pools: Biomass and surface soils (Elana LL et al., 2016; TAM Pugh et al., 2016). Surface soil carbon, derived from plant debris and roots, extends the biomass carbon pool. However, both types of carbon pools have a finite capacity for storage and cannot expand indefinitely (Fang JY, 2021; Piao SL et al., 2022; Zhao HX et al., 2024).

The soil carbon pool, the largest within a terrestrial ecosystem, holds 2–3 times more carbon than the biomass pool and thrice the carbon stored in the atmosphere (Tan ZX et al., 2004; Wang HY et al., 2023). Soil respiration releases CO_2, accounting for approximately two-thirds of the total carbon exchange between terrestrial ecosystems and the atmosphere, making the soil carbon pool a crucial component and carrier of the carbon cycle (Wang YX et al., 1999). Consequently, the soil carbon pool's balance is intrinsically linked to atmospheric CO_2 concentration changes, significantly influencing global climate change (Anil CS et al., 2023; Zhang YX et al., 2022).

In addition to emitting CO_2 into the atmosphere, a soil carbon pool serves as a primary carbon source for groundwater (Kartik J et al., 2022; Li L et al., 2022). In other words, groundwater acts as a significant repository for soil carbon, especially the organic carbon from surface soils. Under biochemical reactions, this organic carbon readily transforms into soluble humic acids like humic acid, fulvic acid, and humin (Li YM et al., 2017; Liu YT et al., 2017). These humic acids from the surface soils infiltrate into groundwater in substantial amounts via processes such as

First author: *E-mail address*: 19809730@qq.com (Hong-yun Ma).

* Corresponding author: *E-mail address*: 358686536@qq.com (Jiang Liu).

Literary editor: Li-qiong Jia
doi:10.31035/cg20230102
2096-5192/© 2025 China Geology Editorial Office.

atmospheric precipitation and irrigation, thus becoming the primary carbon source in groundwater. Consequently, the groundwater carbon cycle represents a crucial extension of soil carbon migration.

The questions then arise: What is the share of the groundwater carbon cycle, and does it have the capacity to influence soil carbon storage? In other words, could the groundwater carbon pool act as an adjunct to the soil carbon pool? These aspects warrant further investigation and clarification.

This study, focusing on the Yinchuan Basin, quantified the organic and total carbon in the surface soil pool, examined the groundwater carbon cycle process, identified the carbon sources in groundwater flow, explored the bases for the evaluation of the groundwater carbon pool, and ultimately juxtaposed the storage of the groundwater carbon pool with that of the soil carbon pool. This study aims to shed light on the impact of the groundwater carbon cycle on the overall atmospheric carbon cycle.

2. Methods

2.1. Study area

The Yinchuan Basin is situated between longitudes 105°45′ and 106°56′ E and latitudes 37°46′ and 39°23′ N. It stretches along the Yellow River, extending 165 km from north to south and spanning 42 km to 60 km from east to west, covering an area of roughly 6138 km². The basin is flanked by the Ordos Block to the east and the Alashan Block to the west, occurring within a subsided depression at the junction of these two blocks (Huang XF et al., 2016). Its terrain is predominantly flat, with elevations ranging from 1100 m to 1200 m (Fig. 1b).

Characterized by an inland arid climate, the Yinchuan Basin experiences a typical continental climate with long,




Fig. 1. Location of the Yinchuan Basin.

cold winters, short, hot summers, rapid spring warming, and early autumn cooling. This region is noted for its low rainfall, ample sunlight, and intense radiation. The basin exhibits an average annual temperature of 8.92°C and an average annual rainfall of 186.7 mm. Rainfall predominantly occurs from June to September, contributing to 68.1% of the yearly total. The annual potential evaporation is 1838.44 mm, with 93.2% occurring from April to September. The average annual humidity is 55%.

The Yinchuan Basin is a fault basin formed due to the gradual distancing of the Ordos and Alashan blocks since the Cenozoic era. This ongoing separation has led to consistent sinking and deposit accumulation within the basin, with sedimentary layers ranging from 400 m to 1600 m in thickness. The central part of the basin witnesses the densest sediment deposition. The basin's eastern and western sides are characterized by plains dominated by diluvial and alluvial-diluvial deposits, whereas its heart is occupied by plains primarily formed from lacustrine deposits, as depicted in the profile in Fig. 1c.

The sedimentary strata within the Yinchuan Basin have created an efficient groundwater aquifer system, which is primarily composed of sand, known for its excellent water retention and conductivity. The lacustrine environments contribute fine-grained clay layers, typically in lens shapes, which do not form a continuous aquitard layer. As a result, groundwaters at varying depths are hydraulically interconnected, establishing a vast and thick aquifer system. This interconnectedness makes it challenging to categorically distinguish between phreatic and confined groundwater. Instead, groundwater is classified based on depth: Shallow groundwater (0−50 m), upper deep groundwater (50−150 m), and lower deep groundwater (150−300 m). Currently, shallow groundwater is primarily exploited for rural domestic use and irrigation. Since 2019, urban domestic water supply has predominantly relied on Yellow River water, with groundwater serving as a secondary, emergency resource.

The Yinchuan Basin's water resources primarily originate from the Yellow River, which serves as the basin's central natural system. Entering from the southwest, the river traverses the plain in a south-north trajectory on the eastern side, covering a distance of 191 km and carrying an annual runoff of approximately 18×10^9 m^3/a. The river's high-quality water supplies the Yinchuan Basin with ample freshwater. Currently, the Yellow River's water is predominantly utilized for agricultural irrigation and some industrial activities, with an average yearly withdrawal of 6.5×10^9 m^3/a, representing roughly a third of the river's total runoff. The basin's other streams have minimal flow and poor water quality, often with total dissolved solids (TDS) content ranging from 3 g/L to 5 g/L, and remain largely untapped.

Groundwater in the basin is chiefly recharged by the infiltration of irrigation water. The Yinchuan Basin boasts an extensive drainage and irrigation infrastructure, including six major canals for irrigation and nine main drainage ditches, complemented by numerous subsidiary branches. This network functions akin to the pulmonary arterial and venous systems, distributing vast quantities of Yellow River water across the basin and then recollecting and redirecting the drainage back to the river. This facilitates a rapid and voluminous groundwater turnover, ensuring efficient water circulation vital for the carbon cycle.

2.2. Data sources

To calculate the carbon storage of surface soils within the Yinchuan Basin, it is essential to obtain data on the carbon content of the surface soil layer and the land use status in the basin.

The carbon contents in the surface soils in the Yinchuan Basin were derived from the project entitled *Multi-Purpose Regional Geochemical Survey of the Yinchuan Basin in the Ningxia Hui Autonomous Region*, conducted by the Department of Natural Resources of Ningxia Hui Autonomous Region in 2007 (Li TB et al., 2012). This project collected a total of 4796 samples, including 97 from construction land areas encompassing urban construction and planning areas and rural homesteads, 1673 from farmland areas such as paddy fields and arid land, 2637 from grassland areas including artificial, natural, and marsh grasslands, 140 from forestland areas including shrubs, trees, and orchards, and 249 from barren land areas like sandy land, bare land, inland beaches, and bare rock land. The statistics of these samples are summarized in Table 1. The variability of data of each category was indicated by the coefficient of variation (CV), the ratio of the standard deviation to the mean of the data. The construction and sandy lands exhibit high CV values, indicating significant variation in their carbon content. In contrast, farmland, grassland, and forestland displayed lower CV values, suggesting more uniform carbon contents

Table 1. Statistics of organic carbon and total carbon contents in the different land use types of the Yinchuan Basin.

Land use type	Samples	Area	Total organic carbon content/%				Total carbon content/%			
		km^2	Average	Max	Min	CV /%	Average	Max	Min	CV /%
Construction land	97	654	0.71	2.96	0.12	74.11	1.88	4.79	0.47	45.10
Farmland	1673	3913	0.60	1.31	0.08	39.70	1.99	3.76	0.18	30.87
Grassland	2637	306	0.30	1.29	0.06	59.22	1.07	2.79	0.09	48.79
Forestland	140	710	0.39	0.84	0.08	49.94	1.41	2.67	0.09	48.93
Barren land	249	159	0.19	0.64	0.05	73.13	0.67	2.40	0.01	89.66
Water area	0	397	–	–	–	–	–	–	–	–
Total	4796	6138	–	–	–	–	–	–	–	–

among their samples. More detailed information is available in the National Geological Archives of China.

The land use data for the Yinchuan Basin were sourced from China's Third National Land Survey (Fig. 1b), covering a total area of 6138 km², including 654 km² of construction land, 3913 km² of farmland (the largest category), 306 km² of grassland, 710 km² of various forestland, 159 km² of barren land (the smallest category), and 397 km² of water area (Table 1). The water area was excluded from the calculation of the surface soil carbon pool.

Calculating the carbon amount involved in the carbon cycle in a groundwater system necessitates the use of groundwater geochemical data of shallow layers. The groundwater geochemical data for the Yinchuan Basin were sourced from projects entitled *A Comprehensive Geological Survey of the Ningxia Yanhuang Economic Zone* and *A Comprehensive Geological Survey of the Ningxia Yanhuang Ecological Economic Zone*, conducted by the Xi'an Center of the China Geological Survey (CGS) from 2016 to 2018 and from 2019 to 2021, respectively. A total of 452 pieces of geochemical data from shallow groundwater (less than 50 m deep) were chosen for analysis.

3. Results and discussion

3.1. Storage assessment of surface soil carbon pools

The soil carbon pool consists of two components: The organic carbon pool and the inorganic carbon pool. Organic carbon in the surface soil primarily originates from plant litter and root systems, with its accumulation in the soils being influenced by the rates of organic matter input, as well as the decomposition and migration rates of soil organic matter (Bao LR et al., 2023). In contrast, inorganic carbon predominantly comprises carbonate minerals like $CaCO_3$, along with dissolved inorganic carbon (HCO_{3-}) in soil moisture and carbon dioxide (CO_2) gas.

Compared to inorganic carbon, organic carbon is more susceptible to microbial breakdown, a complex process that yields various humic acids and eventually releases CO_2. This CO_2 often readily escapes back into the atmosphere, making the link between the organic carbon pool and the atmospheric carbon pool a focal point of concern. Inorganic carbon, however, tends to be more stable.

Since the 1950s, efforts have been made to gauge the carbon stocks of global soil organic carbon pools (Jia YP, 2004). The estimations largely rely on the organic carbon concentration at depths less than 1 m, as soil organic carbon is predominantly found within this top layer. In reality, the concentration of soil organic carbon is especially high in the upper 50 cm, with many soil geochemical surveys concentrating on the top 20 cm. Therefore, a 50 cm depth was chosen as the representative thickness for surface soils in this study.

The soil bulk density in the Yinchuan Basin generally ranges from 1.2 g/L to 1.8 g/L, with an average of 1.5 g/L, which was applied in the assessment of this study.

This study employed the mean contents of organic carbon and total carbon content in samples of each land use to represent the averages of total organic carbon (TOC) and total carbon (TC) in the surface soils of the land type. These values were then multiplied by the total soil mass for a 50 cm depth per unit area and by the total area of the land type. Consequently, the storage of the Yinchuan Basin's surface soil organic carbon pool was estimated at 24.16 Tg and the basin's total carbon storage in surface soil at 61.07 Tg. Detailed calculations are presented in Table 2.

3.2. Analysis of carbon sources in groundwater

Groundwater, as a solution, typically contains seven primary solutes, consisting of four key cations, i.e., Na^+, K+, Ca^{2+}, and Mg^{2+}, and three principal anions, i.e., Cl^-, SO_4^{2-}, and HCO_3^-. In contrast, other elements in the solute composition generally exhibit minor concentrations. These components primarily enter the groundwater system through dissolution, except for HCO_3^-. Analyses indicate that the major anions and cations in groundwater gradually dissolve into it during the groundwater runoff. However, carbon often comes from the soil layer and has already reached a saturation or near-saturation state at the onset of groundwater formation. The subsequent entry of carbon into the groundwater in the runoff process is minimal.

As shown in Fig. 2, the correlation between the total anions (Te^-) and total cations (Te^+) was depicted through a fitted line with a slope of 0.99, demonstrating a strong linear relationship. Excluding SO_4^{2-} and Cl^- separately from the total anions would significantly alter the slopes of the fitted lines displaying the correlation between the remaining anions and the total cations, dropping from 0.99 to 0.62 and 0.40, respectively. The standard deviation (R^2) for the comparative data in each dataset exceeded 0.9, indicating a robust correlation between the data. The changes in these anion content positively correlated with a shift in the total solute concentration in groundwater, suggesting that SO_4^{2-} and Cl^- progressively enter the groundwater system through dissolution as groundwater flows.

Similarly, when comparing total cations to total anions in groundwater (Fig. 3), removing Ca^{2+} and Mg^{2+} or Na^+ and K^+ from the cations and then comparing with the total anions revealed a significant decline in the slopes of the fitted lines, from 1 to 0.69 or 0.31, respectively. This suggests that all

Table 2. Calculated organic carbon and total carbon storage in the various land use types of Yinchuan Basin.

Land use type	Area	TOC content	TC content	TOC	TC
	km²	%	%	Tg	Tg
Construction land	654	0.71	1.88	3.49	14.54
Farmland	3913	0.60	1.99	17.70	38.34
Grassland	306	0.30	1.07	0.69	2.96
Forestland	710	0.39	1.41	2.05	4.47
Barren land	159	0.19	0.67	0.23	0.76
Water area	397	–	–	–	–
Total	6138	–	–	24.16	61.07

major cations, sulfate ions, and chloride ions exhibited similar trends and positively correlated with the total concentration of groundwater, meaning that they all gradually dissolve into the groundwater during the groundwater runoff.

However, the HCO_3^- concentration changed in a significantly different way from those of other ions. As shown in Fig. 2, excluding HCO_3^- led to insignificant shifts in the slope of the fitted line representing the relationship between remaining anions and total cations, only reducing from 0.99 to 0.95. This implies that the HCO_3^- concentration roughly follows the same trend as the total concentration increases. Rather than gradually dissolving into groundwater during the groundwater runoff, carbon has already attained a certain concentration during groundwater formation, with no substantial changes thereafter.

Fig. 4 illustrates the comparison between the total anions and total dissolved carbon in groundwater, which included carbon in various hydrochemical forms such as HCO_3^-, CO_3^{2-}, and free CO_2. The total dissolved carbon concentration and the total anion concentration exhibited a weak correlation, as evidenced by an R^2 value of 0.36.

As depicted in Fig. 5, the ratio of total dissolved carbon to total anions (TC/Te^-) declined with an increase in the total anion concentration. This trend suggests that the dissolution of Cl^- and SO_4^{2-} during the groundwater runoff progressively diminishes the proportion of carbon ions to total anions. In conclusion, carbon does not gradually dissolve and enter the groundwater during the groundwater runoff, making a minimal contribution to the increasing total anion concentration.

Theoretically, given that HCO_3^- represents the primary form of carbon in groundwater, its concentration should align with that of Ca^{2+} if the carbon present in groundwater primarily results from the dissolution of $CaCO_3$.

Nevertheless, a comparison between the Ca^{2+} concentration and the total dissolved carbon concentration in groundwater (Fig. 6) revealed a relatively weak positive correlation, evidenced by a modest slope of 0.26 and an R^2 value of only 0.35. This suggests that the carbon found in the Yinchuan Basin's groundwater might not predominantly originate from $CaCO_3$ dissolution. Instead, it is more likely that the carbon derives from the infiltration of soluble humic acids formed by organic carbon decomposition, though




Fig. 2. Anion concentration *vs.* total cation concentration in groundwater.




Fig. 3. Cation concentration *vs.* total anion concentration in groundwater.




Fig. 4. Total anion concentration *vs.* total dissolved carbon concentration in groundwater.




Fig. 5. Total anion concentration *vs.* ratio of total dissolved carbon to total anions in groundwater.




Fig. 6. Calcium concentration *vs.* total dissolved carbon concentration in groundwater.

conclusive evidence for this is lacking.

In conclusion, the presence of carbon in groundwater is primarily attributed to infiltration from the surface soil carbon pools rather than from a gradual dissolution during groundwater runoff. Thus, groundwater serves as a significant outlet for the surface soil carbon pool.

Further exploration is required to determine the reasonable range for the groundwater carbon pool as a downstream recipient of the surface soil carbon pool.

3.3. Storage assessment of groundwater carbon pool

The Yinchuan Basin boasts an extensive groundwater system, with the Quaternary aquifer's thickness generally ranging from 400 m to 1600 m, supporting various levels of groundwater circulation. The shallow groundwater circulates at depths up to 50 m, with a cycle lasting several decades, whereas deep groundwater circulation extends down to 300 m, with cycles spanning thousands or even tens of thousands of years.

Strictly speaking, all carbon present in groundwater should be considered part of the groundwater carbon pool, which may be extremely large (Zhang SJ et al., 2022; Zhu LL et al., 2022). However, quantifying this is arguably impractical since most groundwater carbon serves merely as a reserve, not interacting with the atmosphere over relevant timescales (beyond a century). The focus should instead be on the portion of groundwater carbon storage that maintains a close connection with the surface soil carbon pool or the atmospheric carbon pool. This portion should have the potential to return to the surface within a specific timeframe and engage in carbon exchange with the atmosphere.

The Yinchuan Basin possesses a comprehensive drainage and irrigation network, with drainage ditches serving as the primary outlet for shallow groundwater. The volume of water discharged through these ditches is substantial, amounting to 18×10^8 m^3/a. This volume significantly surpasses other forms of groundwater discharge, such as manual extraction and runoff to the Yellow River, which are 2 to 3 orders of magnitude smaller and thus negligible for this discussion. Furthermore, the majority of the water in the drainage ditches originates from shallow groundwater discharge. In contrast, contributions from other sources like precipitation, urban domestic drainage, and lake runoff are 2 to 3 orders of magnitude smaller, thereby not influencing the ditches' water volume. Consequently, the water volume in the drainage ditches remains relatively constant over the years, making it a reliable indicator of the active groundwater circulation volume in the Yinchuan Basin.

The active groundwater circulation cycle, also reflected in the drainage water age within the ditches, was assessed using the tritium abundance in various water bodies across the basin (Fig. 7; Ma HY et al., 2019). The cycle period of shallow groundwater was determined based on the average tritium abundance (^{3}H = 53.67 TU) in the basin's main water supply (the Yellow River) and that in the primary discharge (drainage water for irrigation), which averages ^{3}H = 7.89 TU, yielding a calculated cycle period of 33.6 years. The total active groundwater circulation volume was estimated at approximately 604.8×10^8 m^3, derived by multiplying the average annual drainage water volume (18×10^8 m^3/a) by the cycle period (33.6 years).

The average cyclic volume, calculated by dividing the total active groundwater circulation volume (604.8×10^8 m^3) by the total area of the Yinchuan Basin (6138 km²), is 0.985×10^6 m^3/km². The cyclic depth, derived from the average cyclic volume and the average porosity of the shallow aquifer (0.23), is 42.8 m. When considering aquitards, which may act as impermeable barriers within the shallow aquifer and thus do not contribute to active groundwater circulation, the cyclic depth is highly consistent with the commonly accepted shallow groundwater circulation depth of approximately 50 m in the region.

Considering the active groundwater circulation volume in the Yinchuan Basin, which is 604.8×10^8 m^3, and the average total dissolved carbon in the shallow groundwater, measured at 81.01 mg/L derived from 452 samples, the carbon storage in the groundwater pool was estimated at approximately 4.90 Tg (expressed as carbon). This represents 20.27% of the total organic carbon in the surface soils (24.16 Tg) and 8.02% of the total carbon in the surface soils (61.07 Tg). Compared to




Fig. 7. Tritium abundance in different water bodies across the Yinchuan Basin (error lines denoting the distribution ranges of tritium).

the surface soil carbon pool, the carbon storage in the shallow groundwater pool of the Yinchuan Basin is significant and should not be overlooked.

3.4. Doubtful issues deserving discussion

Evaluating the storage of the surface soil carbon pool entails some uncertainties. The first uncertainty is the soil layer thickness of the Yinchuan Basin, which varies significantly across different regions, ranging from 10 cm to 70−90 cm, making it challenging to adjust the assessment on a basin-wide scale to the specific depth of each soil layer. Furthermore, sampling and testing targeted the layers at depths ranging from 0 cm to 20 cm or from 0 cm to 50 cm, with fewer samples taken from depths greater than 50 cm. However, the carbon content is not uniform across depths, tending to decrease sharply with increasing soil depth. Consequently, sample results may not accurately reflect the carbon content across various soil layers, introducing potential errors in the assessment. Some evaluations of surface soil carbon pools' storage consider a depth of 1 m. However, the lack of data on soil layers in this study has hampered a detailed understanding of the vertical distribution of organic carbon in the Yinchuan Basin. Given this, a 50 cm depth was adopted for this assessment. The organic carbon storage in the surface soils of the Ningxia irrigation area with water diverted from the Yellow River, as calculated by the Institute of Soil Science, Chinese Academy of Sciences, is approximately 17.65 Tg (Dong LL et al., 2015), closely aligning with the current estimate of 17.70 Tg. This congruence suggests that the chosen assessment depth is appropriate.

The second uncertainty centers on defining the extent of the groundwater carbon pool. In this study, attention was directed toward shallow groundwater, the most dynamically circulating water in the Yinchuan Basin, while the contributions of middle and deep groundwater were overlooked. This approach is justified by the lengthy cycle periods of deeper groundwater layers, whose carbon content is minimally connected to the atmospheric and surface soil carbon pools. Consequently, the authors agree to primarily consider the actively circulating groundwater for assessments of the groundwater carbon pool, with the definition of actively circulating groundwater being subject to the unique conditions of different watersheds. In this study, the annual discharge from drainage ditches was used to represent active groundwater circulation, deemed to be a suitable metric for evaluation.

The study area's distinct characteristics, particularly its reliance on flood irrigation for agriculture, resulting in substantial irrigation and drainage volumes, significantly enhance the circulation rate of shallow groundwater. This, in turn, contributes to the notable size of the groundwater carbon pool.

When these aspects are set aside and the volumes are compared, the storage estimations of the groundwater carbon pool and the surface soil organic carbon pool are found to have similar orders of magnitude. Thus, the groundwater carbon pool is substantial and warrants increased attention.

4. Conclusions

Evaluations and comparisons between the surface soil carbon pool and the groundwater carbon pool in the Yinchuan Basin lead to several conclusions:

(i) At a 50 cm depth, the surface soils in the Yinchuan Basin hold an organic carbon storage of 24.16 Tg and a total carbon storage of 60.01 Tg.

(ii) Groundwater geochemical characteristics suggest that carbon in groundwater predominantly originates from the infiltration process. Thus, the migration of groundwater carbon represents a significant outflow from the surface soil carbon pool.

(iii) The most dynamically active groundwater in the Yinchuan Basin is the shallow groundwater (less than 50 m deep), which can be considered representative of the groundwater carbon pool's storage. This storage amounts to 4.90 Tg, a substantial figure that cannot be overlooked when compared to the surface soil organic carbon pool, accounting for 20.3%.

Accordingly, it is recommended that the groundwater carbon pool be regarded as an ancillary carbon pool to the surface soils, warranting concurrent evaluation and considerable focus. This approach will enhance the precision of terrestrial carbon balance studies and provide a robust scientific foundation for more effectively meeting the dual carbon goals.

CRediT authorship contribution statement

Hong-yun Ma conceived of the presented idea and performed the computations. Hong-na Ma assisted in plotting the maps. All authors discussed the results and contributed to the final manuscript.

Declaration of competing interest

The authors declare no conflicts of interest.

Acknowledgment

This research was jointly supported by the third scientific survey project in Xinjiang (2022xjkk0300) and the public welfare geological survey projects initiated by the China Geological Survey (DD20190296, DD20221731).

References

Anil CS, Javid ML, Vijayasatya NC, Girisha KG, Olabiyi O, Jeff AB. 2023. Response of soil microbial Communities, inorganic and organic soil carbon pools in arid saline soils to alternative land use practices. Ecological Indicators, 150, 110227. doi: 10.1016/j.ecolind.2023.110227.

Bao LR, Li Y, Zang FL, Yue PJ, Luo K, Deng H, Yan SM. 2023. Spatial distribution, influencing factors and contribution to carbon emission

reduction of topsoil organic carbon in Chongqing, China. Geology in China, https://kns.cnki.net/kcms2/detail/11.1167.P.20230731.1716.004.html (in Chinese with English abstract).

Dong LL, Yu DS, Zhang HD, Shi XZ. 2015. Temporal and spatial changes of soil organic carbon density in Ningxia irrigation zone from 1980 to 2009. Chinese Journal of Ecology, 34(8), 2245–2254 (in Chinese with English abstract). doi: 10.13292/j.1000-4890.2015.0190.

Elana LL, Neal EB, Karl WW. 2016. Source-to-sink sedimentary systems and global carbon burial: A river runs through it. Earth-Science Reviews, 153, 30–42. doi: 10.1016/j.earscirev.2015.10.011.

Fang JY. 2021. Ecological perspectives of carbon neutrality. Chinese Journal of Plant Ecology, 45(1), 1173–1176 (in Chinese with English abstract). doi: 10.17521/cjpe.2021.0394.

Huang XF, Feng SY, Gao R, Li WH. 2016. Development of the Yinchuan Basin: Deep seismic reflection profile revealed the linkages between shallow geology and deep structures. Chinese Journal of Geology, 51(1), 53–66 (in Chinese with English abstract). doi: 10.3969/j.issn.0563-5020.2016.01.006.

Jia YP. 2004. Study advance in the distribution and storage of soil carbon. Journal of Taiyuan Teachers College (Natural Science Edition), 3(4), 62–64 (in Chinese with English abstract).

Kartik J, Cole JC, Kevin GM, K. Ulrich M. 2022. Modeling evaluation of physiochemical processes controlling gas migration in shallow groundwater systems. Advances in Water Resources, 171, 104362. doi: 10.1016/j.advwatres.2022.104362.

Li L, Wu YH, Chong S, Wen QZ, Lin QG, Zhang S. 2022. Influence of CO_2 migration from geological storage on the chemical composition of groundwater and monitoring indicators. Environmental Earth Sciences, 81, 1866–6299. doi: 10.1007/s12665-022-10194-2.

Li TB, Li BG, Gao Y. 2012. A report on the multi target regional geochemical survey of the Yinchuan Basin, Ningxia (scale: 1∶250000). Ningxia Land and Resources Survey and Monitoring Institute, National Geological Data Center, China, 2021-04-22.10.35080/n01. c. 119458 (in Chinese).

Li YM, Gong L, Xie LN. 2017. Soil organic carbon content and carbon mineralization characteristics under different land use types in Northern Tarim Basin. Bulletin of Soil and Water Conservation, 37(3), 216−221 (in Chinese with English abstract). doi: 10.13961/j.cnki.stbctb.2017.03.036.

Liu YT, Gong L, Liu ZY. 2017. Organic carbon and carbon mineralization characteristics under different soil types in southern edge of Tarim Basin. Journal of Arid Land Resources and Environment, 31(2), 162–166 (in Chinese with English abstract). doi: 10.13448/j.cnki.jalre.2017.061.

Ma HY, Li CZ, Wang HQ, Guo L. 2019. Hydrogen and oxygen isotopic compositions of water bodies at Yinchuan Basin. Northwestern Geology, 52(2), 218–225 (in Chinese with English abstract). doi: 10.19751/j.cnki.61-1149/p.2019.02.023.

Piao SL, Zhang XP, Chen AP, Liu Q, Lian X, Wang XH, Peng SS, Wu XC. 2019. The impacts of climate extremes on the terrestrial carbon cycle: A review. Science China Earth Sciences, 62(10), 1551–1563. doi: 10.1007/s11430-018-9363-5.

Piao SL, He Y, Wang XH, Chen FH. 2022. Estimation of China's terrestrial ecosystem carbon sink: Methods, progress and prospects. Earth Sciences, 65(4), 641–651. doi: 10.1007/s11430-021-9892-6.

Tan ZX, R Lal, NE Smeck, FG Calhoun. 2004. Relationships between surface soil organic carbon pool and site variables. Geoderma, 121, 187–195. doi: 10.1016/j.geoderma.2003.11.003.

TAM Pugh, C Müller, A Arneth, V Haverd, B Smith. 2016. Key knowledge and data gaps in modelling the influence of CO^2 concentration on the terrestrial carbon sink. Journal of Plant Physiology, 203, 3–15. doi: 10.1016/j.jplph.2016.05.001.

Wang HY, Wu JQ, Li G, Yan LJ, Liu SN. 2023. Effects of extreme rainfall frequency on soil organic carbon fractions and carbon pool in a wet meadow on the Qinghai-Tibet Plateau. Ecological Indicators, 146, 109853. doi: 10.1016/j.ecolind.2022.109853.

Wang K, Piao SL, He Y, Liu YW, He HL. 2023. Spatial variations and mechanisms for the stability of terrestrial carbon sink in China. Earth Sciences, 66(2), 227–236. doi: 10.1007/s11430-021-1003-5.

Wang YX, Zhao SD, Niu D. 1999. Research state of soil carbon cycling in terrestrial ecosystem. Chinese Journal of Ecology, 18(5), 29–35 (in Chinese with English abstract). doi: 10.13292/j.1000-4890.1999.0070.

Zhang YX, Tang JR, Niu YZ, Zhang JY, Zhao Y, Wei JS, Jiang GZ, Wang LW. 2022. Resource advantage and geological work suggestions under carbon neutralization in Northwest China. Geology in China, 49(5), 1458–1480 (in Chinese with English abstract). doi: 10.12029/gc20220507.

Zhao HX, Fan JD, Gu BJ, Chen YJ. 2024. Carbon sink response of terrestrial vegetation ecosystems in the Yangtze River Delta and its driving mechanism. Journal of Geographical Sciences, 34(1), 112–130. doi: 10.1007/s11442-024-2197-z.

Zhang SJ, Du WF, Liu ZW, Gu XY. 2022. Migration and pollution control of chlorinated hydrocarbons in groundwater system of eastern Jinan. Environmental Earth Sciences, 82, 1866–6299. doi: 10.1007/s12665-022-10691-4.

Zhu LL, Deng ZM, Xie YH, Zhang CY, Chen XR, Li X, Li F, Chen XS, Zou YA, Wang W. 2022. Effects of hydrological environment on litter carbon input into the surface soil organic carbon pool in the Dongting Lake floodplain. CATENA, 208, 105761. doi: 10.1016/j.catena.2021.105761.